

Librería
Bonilla y Asociados
desde 1950



Título: Quadratic Isoperimetric Inequality For Mapping Tori Of Free Group Automorphisms,

Autor: Bridson Martin R/ Groves Daniel

Precio: \$935.61

Editorial:

Año: 2009

Tema:

Edición: 1ª

Sinopsis

ISBN: 9780821846315

The authors prove that if F is a finitely generated free group and ϕ is an automorphism of F then $F \rtimes_{\phi} \mathbb{Z}$ satisfies a quadratic isoperimetric inequality.

The authors' proof of this theorem rests on a direct study of the geometry of van Kampen diagrams over the natural presentations of free-by-cyclic groups. The main focus of this study is on the dynamics of the time flow of t -corridors, where t is the generator of the \mathbb{Z} factor in $F \rtimes_{\phi} \mathbb{Z}$ and a t -corridor is a chain of 2-cells extending across a van Kampen diagram with adjacent 2-cells abutting along an edge labelled t . The authors prove that the length of t -corridors in any least-area diagram is bounded by a constant times the perimeter of the diagram, where the constant depends only on ϕ . The authors' proof that such a constant exists involves a detailed analysis of the ways in which the length of a word $w \in F$ can grow and shrink as one replaces w by a sequence of words w_m , where w_m is obtained from $\phi(w_{m-1})$ by various cancellation processes. In order to make this analysis feasible, the authors develop a refinement of the improved relative train track technology due to Bestvina, Feighn and Handel.

Table of Contents

Positive automorphisms
Train tracks and the beaded decomposition
The General Case
Bibliography
Index