

Librería
Bonilla y Asociados
desde 1950



Título: Kuznetsov's Trace Formula And The Hecke Eigenvalues Of Maass Forms

Autor: A. Knightly

Precio: \$1140.80

Editorial:

Año: 2013

Tema:

Edición: 1^a

Sinopsis

ISBN: 9780821887448

The authors give an adelic treatment of the Kuznetsov trace formula as a relative trace formula on $GL(2)$ over \mathbb{Q} . The result is a variant which incorporates a Hecke eigenvalue in addition to two Fourier coefficients on the spectral side. The authors include a proof of a Weil bound for the generalized twisted Kloosterman sums which arise on the geometric side. As an application, they show that the Hecke eigenvalues of Maass forms at a fixed prime, when weighted as in the Kuznetsov formula, become equidistributed relative to the Sato-Tate measure in the limit as the level goes to infinity.

Table of Contents

!Introduction

!Preliminaries

!Bi-K 8 -invariant functions on $GL(2)(\mathbb{R})$!Maass cusp forms

!Eisenstein series

!The kernel of $R(f)$!A Fourier trace formula for $GL(2)$!Validity of the KTF for a broader class of

h !Kloosterman sums

!Equidistribution of Hecke eigenvalues

!Bibliography

!Notation index

!Subject index