

Librería  
**Bonilla y Asociados**  
desde 1950



**Título:** Kuznetsov's Trace Formula And The Hecke Eigenvalues Of Maass Forms

**Autor:** A. Knightly

**Precio:** \$1140.80

**Editorial:**

**Año:** 2013

**Tema:**

**Edición:** 1<sup>a</sup>

**Sinopsis**

**ISBN:** 9780821887448

The authors give an adelic treatment of the Kuznetsov trace formula as a relative trace formula on  $GL(2)$  over  $\mathbb{Q}$ . The result is a variant which incorporates a Hecke eigenvalue in addition to two Fourier coefficients on the spectral side. The authors include a proof of a Weil bound for the generalized twisted Kloosterman sums which arise on the geometric side. As an application, they show that the Hecke eigenvalues of Maass forms at a fixed prime, when weighted as in the Kuznetsov formula, become equidistributed relative to the Sato-Tate measure in the limit as the level goes to infinity.

Table of Contents

!Introduction

!Preliminaries

!Bi-K 8 -invariant functions on  $GL(2)(\mathbb{R})$  !Maass cusp forms

!Eisenstein series

!The kernel of  $R(f)$  !A Fourier trace formula for  $GL(2)$  !Validity of the KTF for a broader class of  $h$  !Kloosterman sums

!Equidistribution of Hecke eigenvalues

!Bibliography

!Notation index

!Subject index